

Q^2 dependence of chiral-odd twist-3 distribution $e(x, Q^2)$

Y. Koike and N. Nishiyama

Graduate School of Science and Technology, Niigata University, Niigata 950-21, Japan

ABSTRACT

We discuss the Q^2 dependence of the chiral-odd twist-3 distribution $e(x, Q^2)$. The anomalous dimension matrix for the corresponding twist-3 operators is calculated in the one-loop level. This study completes the calculation of the anomalous dimension matrices for all the twist-3 distributions together with the known results for the other twist-3 distributions $g_2(x, Q^2)$ and $h_L(x, Q^2)$. We also have confirmed that in the large N_c limit the Q^2 -evolution of $e(x, Q^2)$ is wholly governed by the lowest eigenvalue of the anomalous dimension matrix which takes a very simple analytic form as in the case of g_2 and h_L .

1. Introduction

The nucleon has three twist-3 distributions g_2 , h_L and e [1]. g_2 is chiral-even and the other two are chiral-odd. e is spin-independent and the other two are spin-dependent. Compared to e , g_2 and h_L have more chance to be measured experimentally since they become leading contribution to proper asymmetries in the polarized deep inelastic scattering (DIS) and Drell-Yan process, respectively. Their Q^2 evolution has been studied in [2,3] for g_2 and in [4,5] for h_L .

Similarly to the distribution functions, there are three twist-3 fragmentation functions which describes hadronization processes of partons in semi-inclusive processes [6], $\hat{g}_2$, $\hat{h}_L$ and $\hat{e}$. (Their naming is parallel to the corresponding distribution functions.) In the inclusive pion production in the transversely polarized DIS, the chiral-odd fragmentation function $\hat{e}$ of the pion appears as a leading contribution together with h_1 (twist-2) of the nucleon. Although the Q^2 evolution of the twist-2 fragmentation functions is known to be obtained from that of the corresponding distributions in the one-loop level (Gribov-Lipatov reciprocity [7]), no such relation is known for the higher twist fragmentation functions.

In our recent work [8], we have investigated the Q^2 evolution of $e(x, Q^2)$. Theoretically, this completed the calculation of the anomalous dimension matrices of all the twist-3 distributions. Phenomenologically, we expect it will shed light on the Q^2 evolution of $\hat{e}(z, Q^2)$, anticipating the day when their relation is clarified. This talk is a brief summary of [8].

2. Q^2 evolution of $e(x, Q^2)$

The chiral-odd twist-3 distribution function $e(x, Q^2)$ is defined by the relation [1]

$$\int \frac{d\lambda}{2\pi} e^{i\lambda x} \langle P | \bar{\psi}(0) \psi(\lambda n) | Q | P \rangle = 2M e(x, Q^2), \quad (1)$$

where $|P\rangle$ is the nucleon (mass M) state with momentum P . Two light-like vectors p and n defined by the relation, $P = p + \frac{M^2}{2}n$, $p^2 = n^2 = 0$, $p \cdot n = 1$, specify the Lorentz frame of the system. Gauge-link operators are implicit in (1). Taking the moments of (1) with respect to x , one can express the moments of $e(x, Q^2)$ in terms of the nucleon matrix elements of the twist-3 operators $V^{\mu_1 \cdots \mu_n}$ (at $m_q = 0$):

$$\mathcal{M}_n[e(Q^2)] = e_n(Q^2), \quad (2)$$

$$\langle PS | V^{\mu_1 \mu_2 \cdots \mu_n} | PS \rangle = 2e_n M (P^{\mu_1} P^{\mu_2} \cdots P^{\mu_n} - \text{traces}), \quad (3)$$

$$V^{\mu_1 \cdots \mu_n} = \sum_{l=2}^n U_l^{\mu_1 \cdots \mu_n} + E^{\mu_1 \cdots \mu_n}, \quad (4)$$

$$U_l^{\mu_1 \cdots \mu_n} = \frac{1}{2} \mathcal{S}_n \bar{\psi} \sigma^{\alpha \mu_1} i D^{\mu_2} \cdots g G_{\alpha}^{\mu_l} \cdots i D^{\mu_n} \psi - \text{traces}, \quad (5)$$

$$E^{\mu_1 \cdots \mu_n} = \frac{1}{2} \mathcal{S}_n \left[\bar{\psi} i \not{D} \gamma^{\mu_1} i D^{\mu_2} \cdots i D^{\mu_n} \psi + \bar{\psi} \gamma^{\mu_1} i D^{\mu_2} \cdots i D^{\mu_n} i \not{D} \psi \right] - \text{traces}, \quad (6)$$

where $\mathcal{M}_n[e(Q^2)] \equiv \int_{-1}^1 dx x^n e(x, Q^2)$ and the covariant derivative $D_\mu = \partial_\mu - ig A_\mu$ restores the gauge invariance.

U_l contains $G_{\mu\nu}$ explicitly, which indicates that $e(x)$ represents the quark-gluon correlations in the nucleon. E is the EOM (equation of motion) operator which vanishes by use of the QCD equation-of-motion. Although the physical matrix elements of EOM operators vanish, one needs to take into account the mixing with E to carry out the renormalization of U_l , as is discussed in [2,4] in the context of the renormalization of g_2 and h_L . From (4) one sees that U_l appears in the form of

$$R_{n,l}^{\mu_1 \cdots \mu_n} = U_{n-l+2}^{\mu_1 \cdots \mu_n} + U_l^{\mu_1 \cdots \mu_n}, \quad \left(l = 2, \dots, \left[\frac{n}{2} \right] + 1 \right). \quad (7)$$

By this combination, $R_{n,l}$ has a definite charge conjugation $(-1)^n$. Note the similarity between the present $\{U_l, R_{n,l}\}$ and $\{\theta_l, R_{n,l}\}$ which appeared in h_L [1,4]. In fact the presence of γ_5 in θ_l is the mere difference from U_l . $R_{n,l}$ of h_L is defined as $\theta_{n-l+2} - \theta_l$ and has a charge conjugation $(-1)^{n+1}$ which is opposite to the above $R_{n,l}$ in (7). We also recall that $e(x)$ does not mix with the gluon-distribution owing to the chiral-odd nature.

For the renormalization of $e(x, Q^2)$, we choose $R_{n,l}$ ($l = 2, \dots, \left[\frac{n}{2} \right] + 1$), E as a basis of the operators. As in the case of g_2 and h_L , We eventually obtained the renormalization constants Z_{ij} among $R_{n,l}$, N and E in the following matrix form:

$$\begin{pmatrix} R_{n,l}^B \\ E_n^B \end{pmatrix} = \begin{pmatrix} Z_{lm}(\mu) & Z_{lE}(\mu) \\ 0 & Z_{EE}(\mu) \end{pmatrix} \begin{pmatrix} R_{n,m}(\mu) \\ E_n(\mu) \end{pmatrix}, \quad \left(l, m = 2, \dots, \left[\frac{n}{2} \right] + 1 \right). \quad (8)$$

This Z_{ij} gives the anomalous dimension matrix for $\{R_{n,l}, E\}$.

3. Large- N_c limit

In [3,5], it has been proved that all the (nonsinglet) twist-3 distributions g_2 , h_L and e obey a simple GLAP equation similarly to the twist-2 distributions in the $N_c \rightarrow \infty$ limit. In this limit, Q^2 evolution of these distributions is completely determined by the lowest eigenvalue of the anomalous dimension matrix which has a simple analytic form. For e , this can be also checked using the obtained result for the mixing matrix Z_{ij} . At $N_c \rightarrow \infty$, i.e., $C_F \rightarrow N_c/2$, the lowest eigenvalue of the anomalous dimension matrix for $R_{n,l}$ is given by (ignoring the factor $g^2/8\pi^2$)

$$\gamma_n^e = 2N_c \left(S_n - \frac{1}{4} - \frac{1}{2(n+1)} \right). \quad (9)$$

Furthermore the Q^2 evolution of e becomes

$$\mathcal{M}_n [e(Q^2)] = L^{\gamma_n^e/b_0} \mathcal{M}_n [e(\mu^2)]. \quad (10)$$

As an example we compare the results for $n = 4$. The exact Q^2 evolution is

$$\begin{aligned} \mathcal{M}_4 [e(Q^2)] &= (0.983a_{4,2}(\mu) + 0.520a_{4,3}(\mu)) \left(\frac{\alpha(Q^2)}{\alpha(\mu^2)} \right)^{9.59/b_0} \\ &+ (0.017a_{4,2}(\mu) - 0.020a_{4,3}(\mu)) \left(\frac{\alpha(Q^2)}{\alpha(\mu^2)} \right)^{15.3/b_0}, \end{aligned} \quad (11)$$

and the leading N_c evolution is

$$\mathcal{M}_4 [e(Q^2)] = \left(a_{4,2}(\mu) + \frac{1}{2}a_{4,3}(\mu) \right) \left(\frac{\alpha(Q^2)}{\alpha(\mu^2)} \right)^{10.4/b_0}, \quad (12)$$

where $a_{n,l}$ is defined by

$$\langle P | R_{n,l}^{\mu_1 \dots \mu_n}(\mu^2) | P \rangle = 2a_{n,l}(\mu^2) M \mathcal{S}_n(P^{\mu_1} \dots P^{\mu_n} - \text{traces}), \quad (13)$$

and $b_0 = \frac{11}{3}N_c - \frac{2}{3}N_f$. One sees clearly that the coefficients in the second term of (11) are small (i.e., $1/N_c^2$ suppressed) and the anomalous dimension in (12) is close to the smaller one in (11).

References

- [1] R.L. Jaffe and X. Ji, Nucl. Phys. **B375**, 527 (1992).
- [2] J. Kodaira, K. Tanaka, T. Uematsu and Y. Yasui, hep-ph 9603377, and references therein.
- [3] A. Ali, V.M. Braun and G. Hiller, Phys. Lett. **B266**, 117 (1991).
- [4] Y. Koike and K. Tanaka, Phys. Rev. **D51**, 6125 (1995).
- [5] I.I. Balitsky, V.M. Braun, Y. Koike and K. Tanaka, hep-ph/9605439, Phys. Rev. Lett. in press.
- [6] R.L. Jaffe and X. Ji, Phys. Rev. Lett. **71**, 2457 (1993).
- [7] V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. **15**, 675 (1972).
- [8] Y. Koike and N. Nishiyama, hep-ph/9609207.